
*

*

// :

// :

			:
			:
			:
/ ±	()	:	
()			
	/ ± / mg/dL	/ ± / mg/dL	:
			:
	/	/	:

()

(counter-regulatory)

± /

()

()

()

()

.()

RAXT

test

[illegible] $/ \pm$

()

.()

-
-

	()	()		
		/		
		/		
		/		
		/		

() () ()

()

.()

.()

()

.()

/ ± / mg/dL

/ ± / mg/dL

.()

()

.() /

.()

%

1. Scott TG. The Effect of Muslim fast of Ramadan on Routine Laboratory Investigation. King Abdulaziz Med J 1981; 1:23-35.

2. Soulaïman AN, Elkhtib FA. Effects of Fasting During Ramadan on Weight and some Blood Constituents in Healthy Muslims. In: The First International Congress on Health and Ramadan. Casablanca, Morocco, Jan 1994, p 16.

3. Nomani MZA, Hallak MH, et al. Changes in Blood Glucose and their Association with Energy-Containing Nutrients in Men on Hypocaloric Diets During Ramadan Fasting. Am J Clin Nutr 1989; 49: 1141-5.

4. Khogheer Y, Sulaiman MI, Al-Fayez SF. Ramadan Fasting State of Controls. Ann Saudi Med 1987; 7 (suppl.): 5-6.

5. Davidson JC. Muslims, Ramadan and Diabetes Mellitus. Bri Med J 1979; 2:1511-2.

6. Groop LC, Bonadonna R, et al: Effect of Prolonged Overnight Fasting on Energy Metabolism in Non-insulin Dependent Diabetic and Non-Diabetic Subjects. Acta Endocrin 1990; 123:30-6.

:

. II

. :

8. Latif AH, Gharieba M Y, Al-Qassab H K, Sartawi O M. Changes in Serum Lipids & Electrolytes in Type II Diabetes Mellitus During Ramadan. Saudi Med J 1993; 14 (6):5323-535.

9. Laajam MA. Ramadan Fasting and Non-Insulin-Dependent Diabetes: Effects on Metabolic Control.

East Afr Med J 1990 Oct; 67 (10): 732-6

10. Bekhadir J, El Ghomari H, et al: Muslims with Non-insulin Dependent Diabetes Fasting During Ramadan: Treatment with Glibenclamide. Bri Med J 1993; 307: 292-5.

11. Ebbing RN. Color Coding of Insulins. Bri Med J 1979; 3:333-4.

12. Sulimani RA, Famuyiwa FO, Laajam M. Diabetes Mellitus and Ramadan Fasting. The Need for a Critical Appraisal. Diabet Med 1988; 5:589-91.

13. Belkhadir J, el Ghomari H, Klicher N, Mikou A, Nasciri M, et al. Muslims with Non-insulin-Dependent Diabetes Fasting During Ramadan: Treatment with Glibenclamide. BM J 1993; 307 (6899): 292-5.

14. Barber SG, Fairweather S, Wright AD, Fitzgerald MG, Malins JM. Muslims, Ramadan and Diabetes Mellitus. BMJ 1979; ii: 46-47 (Letter).

15. The Glimepiride in Ramadan (GLIRA) Study Group. The Efficacy and Safety of Glimepiride in the Management of Type 2 Diabetes in Muslim Patients During Ramadan. Diabetes Care 2005; 28: 421-422.

:

..

17. Salti I, Bénard E, Detournay B, Bianchi-Biscay M, Le Brigand C, Voinet C et al. A Population-Based Study of Diabetes and Its Characteristics during the Fasting Month of Ramadan in 13

Countries: Results of the Epidemiology of Diabetes and Ramadan 1422/2001 (EPIDIAR) Study. *Diabetes Care* 2004; 27: 2306–2311.

18. Yarahmadi S, Larijani B, Bastanhagh MH, Pajouhi M, Baradar Jalili R, Zahedi F, et al. Metabolic and Clinical Effects of Ramadan

Fasting in Patients with Type II Diabetes. *J Coll Physicians Surg Pak* 2003; 13: 329–332.

19. Sari R, Balci MK, Akbas SH, Avcı B. The Effects of Diet, Sulphonylurea and Repaglinide Therapy on Clinical and Metabolic Parameters in

Type 2 Diabetic Patients During Ramadan. *Endocr Res* 2004; 30: 169–177.

20. Uysal AR, Erdogan MF, Sahin G, Kamel N, Erdogan G. Clinical and Metabolic Effects of Fasting in 41 Type 2 Diabetic Patients During Ramadan. *Diabetes Care* 1998; 21:2033–34 (Letter).

21. Salman H, Abdallah MA, Abanamy MA, Al Howasi M. Ramadan Fasting in Diabetic Children in Riyadh. *Diabet Med* 1992; 9: 583–584(Letter).

Effect of Fasting on Blood Sugar in NIDDM Patients in Ramazan Holy Month

Kalantary S.(MD.)

Abstract

Introduction: There is controversy about the effect of fasting on fasting blood sugar of diabetic patients. In this study we are going to determine this issue in the patients with type II diabetics (DM2).

Objective: Survey the effect of fasting on blood sugar in NIDDM patients in Ramazan month.

Materials and Methods: In a quasi experimental study, 17 NIDDM patients (5 males & 12 females) aged 32-65 years old, with the mean of 48.6 ± 8 years old were selected by easy sampling. All patients were on oral agents or diet and fasted during a Ramazan month. Patients with symptomatic diabetic complications & those under 30 & over 65 years old were of this study. Blood was drawn for FBS before & during fasting month at 2 weeks interval. The cases were emphasized to observe the same diet constituents & calorie as before the holy month.

Results: The mean of FBS before & during the holy month was 152 ± 56 mg/dl and 140 ± 25 mg/dl respectively. The difference between fasting blood sugar was not significant.

Conclusion: Fasting did not have any significant effect on FBS in diabetic group of patients.

Key words: Blood Glucose/ Diabetes Mellitus, Insulin- Dependent/ Fasting